

REPROBIOL SPRL

Améliorer la réponse de superovulation chez les vaches lors d'un traitement avec de la FSH (Stimufol) peut se faire par différentes approches basées sur des ajustements de la posologie, la synchronisation du cycle, et la combinaison avec d'autres hormones. Voici quelques conclusions tirées des recherches pertinentes :

1. L'administration d'une faible dose de FSH au début du cycle œstral peut augmenter le taux d'ovulation et la récupération des embryons chez les vaches non-lactantes ([Rajamahendran et al., 1987](#)).

Effect of low dose of FSH given at the beginning of the estrous cycle and subsequent superovulatory response in Holstein cows ☆

R. Rajamahendran¹, R.S. Canseco, C.J. Denbow, F.C. Gwazdauskas², W.E. Vinson

Show more ▾

+ Add to Mendeley ↻ Share 📄 Cite

[https://doi.org/10.1016/0093-691X\(87\)90186-5](https://doi.org/10.1016/0093-691X(87)90186-5)

[Get rights and content](#)

Abstract

A total of 47 superovulations were conducted on forty non-lactating cows to evaluate two different schemes using follicle stimulating hormone (FSH) for superovulating cattle. Cows randomly assigned to treatment A (26 collections) were superovulated beginning on days 9 to 13 of the estrous cycle by giving FSH at decreasing doses of 6, 6, 5, 5, 3, 3, and 2, 2 mg for 4 consecutive days at 12-h intervals while those in treatment B (21 collections) also received 2.5 mg of FSH on days 3 and 4 of the estrous cycle. Animals in both treatments were each given 12.5 mg of prostaglandin $F_{2\alpha}$ ($PGF_{2\alpha}$) at 60 and 72 h after the initiation of superovulatory treatment. Cows were artificially inseminated at 0, 12, and 24 h after the onset of estrus. Embryos were recovered nonsurgically on d 6 and morphologically evaluated. Ovaries of the cows were palpated at the end of flushings to assess the number of corpora lutea (CL). The mean interval from $PGF_{2\alpha}$ to the onset of estrus was not different ($P>0.05$) for treatments A (56.6 h) and B (50.0 h). Also, mean duration of standing estrus was not different for either treatment (13.4 h vs 12.8 h). The mean number of CL palpated (7.3 vs 12.9) and ova recovered (5.5 vs 14.2) were significantly greater ($P<0.05$) for treatment B. The mean number of excellent and good embryos recovered was lower for treatment A animals, but not significant ($P>0.05$). Therefore, low doses of FSH given at the beginning of the cycle increased ovulation rate and embryo recovery in non-lactating cows.



Rue Pelé-Bois 20
4590 Ouffet
Belgium
BE0886.862.189

TÉLÉPHONE 32(0) 477 29 70 18
TÉLÉCOPIE 00-32-86 477 333
E-MAIL info@reprobiol.be
SITE WEB www.reprobiol.be

2. La réduction de la fréquence des injections d'extraits de pituitaire (Stimufol®) pendant la superovulation a montré des résultats similaires à un protocole standard, avec une légère amélioration de la qualité des embryons ([Adel et al., 2018](#)).

Home / Archives /

Vol. 75 No. 1 (2018): BULLETIN OF UNIVERSITY OF AGRICULTURAL SCIENCES AND VETERINARY MEDICINE CLUJ-NAPOCA.
VETERINARY MEDICINE

/

Research Articles

Effect of the Reduction in the Frequency of Injections of Pituitary Extracts During a Superovulation on the Embryo Collection Results in Cows in Algeria

Djallal ADEL

Institute of Veterinary Sciences, University of BLIDA

Kamal TOUATI

University of Liège

Rachid KAIDI

Institute of Veterinary Sciences, University of BLIDA

DOI: <https://doi.org/10.15835/buasvmcn-vm:001217>

Keywords: cattle, embryo production, pituitary extracts, superovulation.

Abstract

The aim of this study is to verify the efficacy of a superovulation treatment with 3 injections of pituitary extracts (Stimufol®) 24 hours apart and compare the results to a standard protocol (8 injections 12 hours apart), with a total dose of 500 µg. Twenty-three embryos collections were conducted after two superovulation protocols, the first with 8 injections 12 hours apart and decreasing doses (n = 11, group 1) and the second 3 injections at constant doses 24 hours of interval (n = 12, group 2). The average number of lutea corpora obtained is 9.1 per cow for group 1 and 10.4 in group 2. A total number of 140 embryos were collected with an average of 6.08 per cow. Among the embryos collected, 46 and 56 per cent were transferable respectively in group 1 and 2. A little quality gain was recorded in group 2 with an average transferable embryo of 3.8 against 2.2 for group 1. In our study we observed the same efficacy for both treatments with a little quality improvement (almost one transferable embryo) when reducing the number of injections at the time of superovulation treatment, the same gain minimal in elite cows can have a significant economic impact.






Rue Pelé-Bois 20
4590 Ouffet
Belgium
BE0886.862.189




TÉLÉPHONE 32(0) 477 29 70 18
TÉLÉCOPIE 00-32-86 477 333
E-MAIL info@reprobiol.be
SITE WEB www.reprobiol.be

3. Les polymorphismes dans la région amont du gène du récepteur de la FSH peuvent être associés aux traits de superovulation chez les vaches Holstein chinoises, indiquant que le récepteur de la FSH est un marqueur potentiel pour la réponse à la superovulation ([Yang et al., 2010](#)).

Polymorphisms in the 5' upstream region of the FSH receptor gene, and their association with superovulation traits in Chinese Holstein cows

[Wu-Cai Yang](#)^a, [Shu-Jing Li](#)^{b 1}, [Ke-Qiong Tang](#)^a, [Guo-Hua Hua](#)^a, [Chun-Yan Zhang](#)^a, [Jun-Na Yu](#)^a,
[Li Han](#)^a, [Li-Guo Yang](#)^a  

Show more 

 Add to Mendeley  Share  Cite

<https://doi.org/10.1016/j.anireprosci.2010.02.004>

[Get rights and content](#) 

Abstract

To identify a predictor to forecast superovulation response on the basis of associations between superovulation performance and gene polymorphism, variation in the bovine follicle stimulating hormone receptor (FSHR) gene was investigated using PCR-single-strand conformational (PCR-SSCP) and DNA sequencing. One single nucleotide polymorphism (SNP) of G-278A located in the 5' upstream region of bovine FSHR gene was found in 118 Chinese Holstein cows treated for superovulation. Two SNPs of G-278A (GU253337) and A-320T (rs43676359) were analyzed. In polymorphic locus -278, all cows without superovulation response were mutations with genotypes of CD and DD. Cows with CC genotype had a significant increase in the total number of ova (TNO) ($P < 0.01$), and produced more transferable embryos (NTE) than those of the CD and DD genotypes ($P < 0.01$). At this locus, the additive effects seemed to be highly significant ($P < 0.01$) and allele C was associated with an increase in the TNO and NTE. These results indicated that the FSHR is a potential marker for superovulation response and can be used as a predictor for superovulation in Chinese Holstein cows.



Rue Pelé-Bois 20
4590 Ouffet
Belgium
BE0886.862.189

TÉLÉPHONE 32(0) 477 29 70 18
TÉLÉCOPIE 00-32-86 477 333
E-MAIL info@reprobiol.be
SITE WEB www.reprobiol.be

4. Les niveaux de FSH dans le plasma des vaches superovulées avec des injections quotidiennes de FSH montrent la nécessité d'utiliser deux injections quotidiennes en raison de la demi-vie relativement courte de la FSH ([Demoustier et al., 1988](#)).



Determination of porcine plasma follitropin levels during superovulation treatment in COWS

[M.M. Demoustier](#)¹, [J.-Fr. Beckers](#)², [P. Van Der Zwalmen](#)², [J. Closset](#)³, [J.-L. Gillard](#)¹, [Fr. Ectors](#)²

Show more ▾

+ Add to Mendeley Share Cite

[https://doi.org/10.1016/0093-691X\(88\)90185-9](https://doi.org/10.1016/0093-691X(88)90185-9)

[Get rights and content](#) ↗


Abstract


Porcine follicle stimulating hormone (pFSH) and porcine luteinizing hormone (pLH), are widely used to induce superovulation in cows. An advantage of this treatment is that the LH:FSH ratio can be varied to optimize the growth of the ovarian follicles. However, due to the relatively short half-life of FSH, the superovulatory treatment requires numerous injections.



A performant radioimmunoassay system (sensitivity=0.2 ng/ml plasma) was used to determine plasma pFSH levels in cows that were superovulated with 2 daily injections of 4 Armour Units (A.U.) of pFSH for 4 d. From plasma profiles, the half-life and the disappearance of pFSH were estimated at 5 h and at 10 to 12 h, respectively, confirming the necessity of using two daily injections.

5. La dynamique folliculaire et la réponse superovulatoire chez les vaches traitées avec la FSH-P dans différents états endocriniens montrent que l'environnement endocrinien peut affecter la réponse à la superovulation ([Savio et al., 1991](#)).

Follicular dynamics and superovulatory response in Holstein cows treated with FSH-P in different endocrine states

J.D. Savio¹, H. Bongers¹, M. Drost², M.C. Lucy¹, W.W. Thatcher¹ 

Show more 

+ Add to Mendeley  Share  Cite

[https://doi.org/10.1016/0093-691X\(91\)90303-U](https://doi.org/10.1016/0093-691X(91)90303-U)

[Get rights and content](#) 

Abstract

The effect of follicular and/or endocrine environments on superovulatory response was tested. Eighteen nonlactating Holstein cows were superovulated with 32 mg FSH-P given in decreasing doses at 12-h intervals plus two injections of prostaglandin F2-alpha (25 mg each) on the third day of treatment. Cows were assigned randomly to treatments: T1, superovulatory treatment initiated on estrous cycle Day 10.5; T2, CIDR (intravaginal device containing 1.9 g of progesterone) inserted from Days 3 to 9 and superovulation initiated on Day 6.5; T3, identical to T2 but Buserelin (GnRH agonist) was injected (8 µg, i.m.) on Day 3 at the time of CIDR insertion. Embryos were recovered on Day 7 after the superovulatory estrus. Cows were examined daily by ultrasonography and blood was collected for progesterone and estradiol determinations. Mean diameter of the dominant follicle (frequency of first-wave dominant follicle) at the beginning of FSH injections was 13.7 mm (46), 11.2 mm (66) and 8.7 mm (66) ($P<0.01$) for T1, T2 and T3, respectively. Following initiation of superovulation, follicles moved into larger follicle classes (Class I, <3 mm; Class II, 3 to 4 mm; Class III, 5 to 9 mm; Class IV >9 mm) earliest in T1 ($P<0.01$). Cumulative follicular diameter and plasma concentrations of estradiol at Day 4 of superovulation were higher ($P<0.01$) in T1 (200 mm, 82 pg/ml) compared with T2 (123 mm, 24 pg/ml) and T3 (130 mm, 18 pg/ml). Proportion of cows in estrus prior to 12 h vs 12 to 24 h differed ($P<0.05$) between groups (T1: 5 vs 1; T2: 2 vs 4; T3: 1 vs 5). Mean number of follicles on the last day of superovulation treatment, number of CL and number of embryos plus unfertilized ova recovered were 17.5, 12.2 and 13.3; 13.8, 10 and 8.2 ($P<0.1$) and 8.7, 4.5 and 2.3 ($P<0.05$) for T1, T2 and T3, respectively. The developmental stage of the dominant follicle was associated with not only the number of ovulations, but also the size and peri-estrous concentrations of plasma estradiol associated with the recruited follicles.



Rue Pelé-Bois 20
4590 Ouffet
Belgium
BE0886.862.189

TÉLÉPHONE 32(0) 477 29 70 18
TÉLÉCOPIE 00-32-86 477 333
E-MAIL info@reprobiol.be
SITE WEB www.reprobiol.be

6. Une corrélation négative a été observée entre l'augmentation de la concentration de progestérone après le début du traitement par FSH et le pourcentage d'embryons transférables collectés, suggérant un impact de la progestérone sur la qualité des embryons ([Tamboura et al., 1985](#)).

Superovulation in cows: A relationship between progesterone secretion before ovulation and the quality of embryos

[D. Tamboura](#), [D. Chupin](#), [J. Saumande](#)

Show more ▾

+ Add to Mendeley [Share](#) [Cite](#)

[https://doi.org/10.1016/0378-4320\(85\)90048-X](https://doi.org/10.1016/0378-4320(85)90048-X)

[Get rights and content](#)

Abstract

Superovulation was induced in 28 cyclic cows by treatment with FSH and milk progesterone concentrations were studied from the initiation of treatment to oestrus. A negative correlation ($r = -0.66$; $P < 0.001$) was observed between the increase in progesterone concentration observed 2 days after the beginning of treatment and the percentage of transferable embryos collected. During this sampling period progesterone secretion was not related to ovulation rate and number of embryos.

Ces études suggèrent plusieurs approches pour améliorer la réponse à la superovulation chez les vaches traitées avec Stimufol, y compris l'ajustement des doses et du calendrier de traitement, ainsi que la considération des facteurs génétiques et endocriniens individuels.

7. **Surveillance Ultrasonographique** : Utiliser l'ultrasonographie pour surveiller le développement folliculaire et ajuster le traitement en conséquence.
8. **Gestion du Stress** : Minimiser le stress des animaux pendant le traitement, car le stress peut affecter négativement la réponse à la superovulation.



Rue Pelé-Bois 20
4590 Ouffet
Belgium
BE0886.862.189

TÉLÉPHONE 32(0) 477 29 70 18
TÉLÉCOPIE 00-32-86 477 333
E-MAIL info@reprobiol.be
SITE WEB www.reprobiol.be