

## REPROBIOL SPRL

Pourquoi le traitement FSH-LH ne provoque pas systématiquement une réponse de superovulation chez les vaches :

Le traitement de FSH ne provoque pas systématiquement une superovulation chez les vaches pour plusieurs raisons, dont certaines sont explorées dans des études récentes :

1. **Variabilité de la réponse ovarienne** : La réponse ovarienne à la FSH peut varier considérablement d'une vache à l'autre, influencée par des facteurs tels que l'âge, la race, la condition corporelle, et le stade du cycle estral au moment du traitement (Bó & Mapletoft, 2014).



Rue Pelé-Bois 20  
4590 Ouffet  
Belgium  
BE0886.862.189

TÉLÉPHONE 32(0) 477 29 70 18  
TÉLÉCOPIE 00-32-86 477 333  
E-MAIL info@reprobiol.be  
SITE WEB www.reprobiol.be

2. **Concentration de progestérone** : Une corrélation négative a été observée entre l'augmentation de la concentration de progestérone observée 2 jours après le début du traitement de superovulation et le pourcentage d'embryons transférables collectés ([Tamboura, Chupin, & Saumande, 1985](#)).




## Superovulation in cows: A relationship between progesterone secretion before ovulation and the quality of embryos

[D. Tamboura](#), [D. Chupin](#), [J. Saumande](#)

I.N.R.A. Station de Physiologie de la Reproduction, 37380 Nouzilly France

Accepted 28 June 1984, Available online 2 October 2003.

Show less 

 Add to Mendeley  Share  Cite

[https://doi.org/10.1016/0378-4320\(85\)90048-X](https://doi.org/10.1016/0378-4320(85)90048-X)

[Get rights and content](#) 

### Abstract

Superovulation was induced in 28 cyclic cows by treatment with FSH and milk progesterone concentrations were studied from the initiation of treatment to oestrus. A negative correlation ( $r = -0.66$ ;  $P < 0.001$ ) was observed between the increase in progesterone concentration observed 2 days after the beginning of treatment and the percentage of transferable embryos collected. During this sampling period progesterone secretion was not related to ovulation rate and number of embryos.




Rue Pelé-Bois 20  
4590 Ouffet  
Belgium  
BE0886.862.189



TÉLÉPHONE 32(0) 477 29 70 18  
TÉLÉCOPIE 00-32-86 477 333  
E-MAIL [info@reprobiol.be](mailto:info@reprobiol.be)  
SITE WEB [www.reprobiol.be](http://www.reprobiol.be)

3. **Demi-vie de la FSH** : En raison de la demi-vie relativement courte de la FSH, le traitement superovulatoire nécessite de nombreuses injections, ce qui peut affecter la réponse ovarienne ([Demoustier et al., 1988](#)).

## Determination of porcine plasma follitropin levels during superovulation treatment in COWS

M.M. Demoustier<sup>1</sup>, J.-Fr. Beckers<sup>2</sup>, P. Van Der Zwalmen<sup>2</sup>, J. Closset<sup>3</sup>, J.-L. Gillard<sup>1</sup>, Fr. Ectors<sup>2</sup>

Show more 

+ Add to Mendeley  Share  Cite

[https://doi.org/10.1016/0093-691X\(88\)90185-9](https://doi.org/10.1016/0093-691X(88)90185-9)

[Get rights and content](#) 

### Abstract

Porcine follicle stimulating hormone (pFSH) and porcine luteinizing hormone (pLH), are widely used to induce superovulation in cows. An advantage of this treatment is that the LH:FSH ratio can be varied to optimize the growth of the ovarian follicles. However, due to the relatively short half-life of FSH, the superovulatory treatment requires numerous injections.

A performant radioimmunoassay system (sensitivity=0.2 ng/ml plasma) was used to determine plasma pFSH levels in cows that were superovulated with 2 daily injections of 4 Armour Units (A.U.) of pFSH for 4 d. From plasma profiles, the half-life and the disappearance of pFSH were estimated at 5 h and at 10 to 12 h, respectively, confirming the necessity of using two daily injections.



Rue Pelé-Bois 20  
4590 Ouffet  
Belgium  
BE0886.862.189

TÉLÉPHONE 32(0) 477 29 70 18  
TÉLÉCOPIE 00-32-86 477 333  
E-MAIL [info@reprobiol.be](mailto:info@reprobiol.be)  
SITE WEB [www.reprobiol.be](http://www.reprobiol.be)

4. **Influence des gonadotrophines endogènes** : Le profil hormonal naturel de la vache, y compris les niveaux de gonadotrophines endogènes comme l'hormone anti-müllérienne (AMH), peut influencer la réponse à la FSH. Les niveaux d'AMH sont corrélés au nombre de follicules antraux et peuvent prédire la réponse ovarienne à la superovulation ([Rico et al., 2009](#)).

**Anti-Müllerian Hormone Is an Endocrine Marker of Ovarian Gonadotropin-Responsive Follicles and Can Help to Predict Superovulatory Responses in the Cow<sup>1</sup>**

Charlène Rico,<sup>3</sup> Stéphane Fabre,<sup>3</sup> Claire Médigue,<sup>5</sup> Nathalie di Clemente,<sup>6</sup> Frédérique Clément,<sup>5</sup> Martine Bontoux,<sup>3</sup> Jean-Luc Touzé,<sup>3</sup> Mickaël Dupont,<sup>4</sup> Eric Briant,<sup>4</sup> Benoît Rémy,<sup>7</sup> Jean-François Beckers,<sup>7</sup> and Danielle Monniaux<sup>2,3</sup>

*Physiologie de la Reproduction et des Comportements, UMR 6175 INRA-CNRS-Université de Tours-Haras Nationaux,<sup>3</sup> and Unité Expérimentale de Physiologie Animale de l'Orfrasière,<sup>4</sup> Centre INRA de Tours, 37380 Nouzilly, France  
Centre de Recherche INRIA Paris-Rocquencourt,<sup>5</sup> Domaine de Voluceau, 78153 Le Chesnay, France  
Endocrinologie et Génétique de la Reproduction et du Développement,<sup>6</sup> Institut National de la Santé et de la Recherche Médicale, U782, Université Paris-Sud, UMR-S0782, F-92140 Clamart, France  
Laboratory of Endocrinology and Animal Reproduction,<sup>7</sup> Faculty of Veterinary Medicine, University of Liege, B-4000 Liege, Belgium*

**ABSTRACT**

The major limitation to the development of embryo production in cattle is the strong between-animal variability in ovulatory response to FSH-induced superovulation, mainly due to differences in ovarian activity at the time of treatment. This study aimed to establish whether anti-Müllerian hormone (AMH) was an endocrine marker of follicular populations in the cow, as in human, and a possible predictor of the ovarian response to superovulation. Anti-Müllerian hormone concentrations in plasma varied 10-fold between cows before treatment and were found to be highly correlated with the numbers of 3- to 7-mm antral follicles detected by ovarian ultrasonography before treatment ( $r = 0.79$ ,  $P < 0.001$ ) and the numbers of ovulations after treatment ( $r = 0.64$ ,  $P < 0.01$ ). Between-animal differences in AMH concentrations were found to be unchanged after a 3-mo delay ( $r = 0.87$ ,  $P < 0.01$ ), indicating that AMH endocrine levels were characteristic of each animal on a long-term period. The population of healthy 3- to 7-mm follicles was the main target of superovulatory treatments, contained the highest AMH concentrations and AMH mRNA levels compared with larger follicles, and contributed importantly to AMH endocrine levels. In conclusion, AMH was found to be a reliable endocrine marker of the population of small antral gonadotropin-responsive follicles in the cow. Moreover, AMH concentrations in the plasma of individuals were indicative of their ability to respond to superovulatory treatments.

*AMH, assisted reproductive technology, follicular development, granulosa cells, ovary*



Rue Pelé-Bois 20  
4590 Ouffet  
Belgium  
BE0886.862.189

TÉLÉPHONE 32(0) 477 29 70 18  
TÉLÉCOPIE 00-32-86 477 333  
E-MAIL info@reprobiol.be  
SITE WEB www.reprobiol.be

5. **Effets du stress et de la manipulation** : Le stress et la manipulation associés au traitement de superovulation peuvent également affecter négativement la réponse des vaches. Des méthodes moins invasives et stressantes pour administrer la FSH pourraient améliorer la réponse ([Sakaguchi et al., 2018](#)).

*Journal of Reproduction and Development, Vol. 64, No 5, 2018*

—Technology Report—

## Effect of a single epidural administration of follicle-stimulating hormone via caudal vertebrae on superstimulation for *in vivo* and *in vitro* embryo production in Japanese black cows

Kenichiro SAKAGUCHI<sup>1, 2)</sup>, Atsushi IDETA<sup>1)</sup>, Yojiro YANAGAWA<sup>3)</sup>, Masashi NAGANO<sup>3)</sup>, Seiji KATAGIRI<sup>3)</sup> and Masato KONISHI<sup>1)</sup>

<sup>1)</sup>Zen-noh Embryo Transfer Center, Hokkaido 080-1407, Japan

<sup>2)</sup>Laboratory of Theriogenology, Graduate School of Veterinary Medicine, Hokkaido University, Hokkaido 060-0818, Japan

<sup>3)</sup>Laboratory of Theriogenology, Department of Clinical Sciences, Faculty of Veterinary Medicine, Hokkaido University, Hokkaido 060-0818, Japan

**Abstract.** Here, we describe a simplified procedure for embryo production in the Japanese black cow that uses a single caudal epidural injection of follicle-stimulating hormone (FSH). First, we compared the efficiency of superovulation for *in vivo* embryo production between conventional multiple FSH treatment (control, n = 10) and single epidural administration (epidural, n = 5). The number of transferable blastocysts was similar between control and epidural groups ( $4.7 \pm 3.5$  and  $9.0 \pm 6.0$ , respectively). Next, we compared *in vitro* embryo production by ovum pick-up and *in vitro* fertilization (OPU-IVF) between control (n = 12) and epidural groups (n = 12). The rate of development to transferable blastocysts was higher in the epidural group than in the control (23.3 vs. 10.5%,  $P < 0.001$ ). In conclusion, a single epidural administration of FSH can induce follicular development comparable to that of the conventional superovulation protocol and may improve the productivity of OPU-IVF.

**Key words:** Epidural administration, Follicle-stimulating hormone (FSH), Ovum-pick up, Superovulation  
(*J. Reprod. Dev.* 64: 451–455, 2018)

6. Une étude a révélé une corrélation négative entre l'augmentation de la concentration de progestérone deux jours après le début du traitement par FSH et le pourcentage d'embryons transférables collectés chez des vaches cycliques. Cette augmentation n'était cependant pas liée au taux d'ovulation et au nombre d'embryons ([Tamboura, Chupin, & Saumande, 1985](#)).

## Superovulation in cows: A relationship between progesterone secretion before ovulation and the quality of embryos

[D. Tamboura, D. Chupin, J. Saumande](#)

Show more 

[+](#) Add to Mendeley [🔗](#) Share [🗒](#) Cite

[https://doi.org/10.1016/0378-4320\(85\)90048-X](https://doi.org/10.1016/0378-4320(85)90048-X)

[Get rights and content](#) 

### Abstract

Superovulation was induced in 28 cyclic cows by treatment with FSH and milk progesterone concentrations were studied from the initiation of treatment to oestrus. A negative correlation ( $r = -0.66$ ;  $P < 0.001$ ) was observed between the increase in progesterone concentration observed 2 days after the beginning of treatment and the percentage of transferable embryos collected. During this sampling period progesterone secretion was not related to ovulation rate and number of embryos.



Rue Pelé-Bois 20  
4590 Ouffet  
Belgium  
BE0886.862.189

TÉLÉPHONE 32(0) 477 29 70 18  
TÉLÉCOPIE 00-32-86 477 333  
E-MAIL [info@reprobiol.be](mailto:info@reprobiol.be)  
SITE WEB [www.reprobiol.be](http://www.reprobiol.be)

7. Une autre étude a examiné l'utilisation de FSH dissoute dans du polyvinylpyrrolidone (PVP) pour réduire le nombre de traitements nécessaires pour induire une superovulation chez les vaches. Les résultats ont suggéré que le PVP est un solvant approprié pour prolonger l'absorption de la FSH administrée en une seule injection, offrant ainsi une approche plus pratique ([Yamamoto, Ooe, Kawaguchi, & Suzuki, 1994](#)).



## Superovulation in the cow with a single intramuscular injection of FSH dissolved in polyvinylpyrrolidone

[M. Yamamoto](#)<sup>2</sup>, [M. Ooe](#)<sup>2</sup>, [M. Kawaguchi](#)<sup>1</sup>, [T. Suzuki](#)<sup>2</sup>

Show more

+ Add to Mendeley Share Cite

[https://doi.org/10.1016/0093-691X\(94\)90184-K](https://doi.org/10.1016/0093-691X(94)90184-K)

[Get rights and content](#)

### Abstract

It is desirable to reduce the number of treatments required to induce superovulation in cows. In this study we examined whether dissolving FSH in polyvinylpyrrolidone (PVP) would reduce the rate of absorption of FSH and allow it to be administered in a single dose for superovulation. In Experiment 1, 10 cows each received a single dose of FSH which contains 0.6% luteinizing hormone (FSH-R; 30mg i.m.) dissolved in 30% PVP (10ml) or in saline. In Experiment 2, a single injection of 30mg FSH-R dissolved in 30% PVP was given to 25 cows, and 32 cows were injected twice daily in declining doses to receive a total of 28mg FSH-R dissolved in saline. Prostaglandin F<sub>2α</sub> was given to all the cows 48h after the first FSH treatment. Embryos were collected on Day 7 or 8 post insemination. In Experiment 1, the effect of FSH dissolved in PVP was compared with that dissolved in saline (number of recovered ova and embryos; 9.4±4.1 vs. 0). In Experiment 2, the rate of transferable embryos by single injection of FSH-R in PVP were significantly higher (P<0.05) than that of treatment of multiple injection groups. Progesterone concentration measured in serum collected 4 times from estrus (Day-0) to the day of embryo collection, indicated similar patterns in the 2 treatment groups. These findings suggest that PVP is a suitable solvent for prolonging the absorption of FSH given in a single injection thus providing a more practical approach of FSH administration.



Rue Pelé-Bois 20  
4590 Ouffet  
Belgium  
BE0886.862.189

TÉLÉPHONE 32(0) 477 29 70 18  
TÉLÉCOPIE 00-32-86 477 333  
E-MAIL [info@reprobiol.be](mailto:info@reprobiol.be)  
SITE WEB [www.reprobiol.be](http://www.reprobiol.be)



8. Une étude a montré qu'une réponse superovulatoire des ovaires chez des vaches buffles a été observée avec l'utilisation de FSH à raison de 40 mg sur quatre jours, suggérant que les protocoles de superovulation peuvent nécessiter des ajustements spécifiques à l'espèce ([Karaivanov et al., 1987](#)).

### [Superovulation induction in buffalo cows (*Bubalus bubalis*) with follicle-stimulating hormone (FSH)].

👤 K. Karaivanov, K. Vlahov, M. Petrov + 2 more authors

📅 1987

#### Abstract

A total of 19 buffalo cows were used to induce superovulation following a pattern that included the use of FSH at the rate of 40 mg in the course of four days, applied twice a day and of Oestrophan at 48 hours after the beginning of treatment. Fifteen animals manifested estrus and were twice inseminated. The superovulation response of the ovaries was 4.3 +/- 0.8 yellow bodies and 0.5 +/- 0.24 follicles greater than 8 mm. A nonsurgical method was employed to obtain 24 embryos of high quality and an unfertilized ovum from 8 buffalo cows.



Rue Pelé-Bois 20  
4590 Ouffet  
Belgium  
BE0886.862.189

TÉLÉPHONE 32(0) 477 29 70 18  
TÉLÉCOPIE 00-32-86 477 333  
E-MAIL [info@reprobiol.be](mailto:info@reprobiol.be)  
SITE WEB [www.reprobiol.be](http://www.reprobiol.be)



9. Les profils hormonaux de FSH, LH, estradiol et progestérone chez les vaches traitées avec une préparation de FSH porcin pour induire une superovulation ont montré des changements significatifs par rapport à un cycle œstral normal, avec une suppression partielle des pics préovulatoires de FSH, bien que les pics de LH n'aient pas toujours été supprimés ([Kaneko et al., 1989](#)).

### Suppression of the Preovulatory Surge of FSH in Superovulating Cattle Pretreated with Porcine FSH

Hiroyuki KANEKO, Takayoshi TERADA, Koichi GOTO, Tatsuhiko NAKASHIMA, Kiyoko OGATA, Masahiro KONDOH, Kazuyoshi TAYA, Shuji SASAMOTO

[+](#) Author information

Keywords: FSH, LH, ESTRADIOL, PROGESTERONE, SUPEROVULATING CATTLE

JOURNAL FREE ACCESS

1989 Volume 35 Issue 1 Pages 7-13

[DOI](#) <https://doi.org/10.1262/jrd1977.35.7>

[+](#) Details

#### Article overview

- > [Abstract](#)
- > [References \(34\)](#)
- > [Content from these authors](#)

#### Share



#### Abstract

Changes in plasma concentrations of FSH, LH, estradiol and progesterone were examined in cows treated with porcine FSH preparation to induce superovulation. The results were compared with those in the normal estrous cycle prior to FSH treatment in the same cows. The number of ovulations in each cow was from 5 to 7. Plasma levels of estradiol before ovulation and of progesterone after ovulation were much higher than those in the normal oestrous cycle. During the period of superovulation, mean concentrations of preovulatory peaks of FSH were suppressed to 61% of normal controls, though, LH peaks were not always suppressed. These results indicate that a partial suppression of the preovulatory FSH surge occurred in cattle when superovulation was induced by exogenous gonadotropin treatment.